

A direct method for measuring heat conductivity in intracluster medium

Makoto Hattori

hattori@astr.tohoku.ac.jp

and Nobuhiro Okabe

okabe@astr.tohoku.ac.jp

*Astronomical Institute, Graduate School of Science, Tohoku University,
Sendai 980-8578, Japan.*

ABSTRACT

The inverse Compton scattering of the cosmic microwave background (CMB) radiation with electrons in the intracluster medium which has a temperature gradient, was examined by the third-order perturbation theory of the Compton scattering. A new type of the spectrum distortion of the CMB was found and named as gradient T Sunyaev-Zel'dovich effect (gradT SZE). The spectrum has an universal shape. The spectrum crosses over zero at 326GHz. The sign of the spectrum depends on the relative direction of the line-of-sight to the direction of the temperature gradient. This unique spectrum shape can be used to detect the gradT SZE signal by broad-band or multi-frequency observations of the SZE. The amplitude of the spectrum distortion does not depend on the electron density and is proportional to the heat conductivity. Therefore, the gradT SZE provides an unique opportunity to measure thermally nonequilibrium electron momentum distribution function when the ICM has a temperature gradient and the heat conductivity in the ICM. However, the expected amplitude of the signal is very small. The modifications to the thermal SZE spectrum due to variety of known effects, such as relativistic correction etc., can become problematic when using multi-frequency separation techniques to detect the gradT SZE signal.

Subject headings: galaxies: clusters: general—magnetic fields—conduction—Compton scattering—plasma—mm and sub-mm observations

1. Introduction

The recent dramatic progresses of the X-ray observations have been unveiling that the temperature distribution of the intracluster medium (hereafter ICM) is far from isothermal. It has turned out that the heat conductivity is one of the key parameters controlling the evolution of the ICM (Ikebe et al. 1999; Markevitch et al. 2000; Fabian et al. 2001; Vikhlinin et al. 2001; Markevitch et al. 2003; Zakamska & Narayan 2003). The global temperature gradient have been found in some cooling core clusters (Kaastra et al. 2004). Zakamska & Narayan (2003) performed hydrostatic model fitting of the observed temperature and density profile of the ICM. They assumed the energy balance between the radiative cooling and the conductive heating. They showed that the heat conductivities should be one-third of the Spitzer value in majority of the clusters. Shock waves (Markevitch et al. 2002; Fujita et al. 2004) and cold fronts (Markevitch et al. 2000; Vikhlinin et al. 2001) found in merging clusters are showing the temperature jumps across the discontinuities. The hot bubbles caused by the interaction with the activity of the central galaxies have been found in many clusters (McNamara et al. 2000; Fabian et al. 2000; Blanton et al. 2001). The heat conductivities across the cold fronts and the surface of the hot bubbles have to be reduced in many order of magnitudes to maintain the structures (Ettori & Fabian 2000). The large scale fluctuation of the temperature distributions have been found in several clusters (Watanabe et al. 1999; Shibata et al. 2001; Markevitch et al. 2003). To maintain the observed temperature fluctuations at least for a dynamical time scale of clusters, the order of magnitude reduction of the heat conductivities are required (Markevitch et al. 2003). All the above mentioned facts show that a direct method for measuring the heat conductivities may provide crucial informations to study the evolution of the ICM.

The recent theoretical progresses have been unveiling that a microscopic instability of the plasma which has a temperature gradient (Ramani & Laval 1978), could play important roles on the suppression of the heat conductivities (Levinson & Eichler 1992; Hattori & Umetsu 2000; Okabe & Hattori 2003) and the origin of cluster magnetic fields (Okabe & Hattori 2003; Okabe & Hattori 2004). The heat conduction is the electron thermal energy flow from hot to cold regions. Since it is the third order moment of the electron momentum distribution function, the electron momentum distribution function must have deviation from the Maxwell-Boltzmann distribution which is the even function of the momentum. Therefore, the electron momentum distribution function of the plasma with a temperature gradient is described by the sum of the Maxwell-Boltzmann distribution, $g_m(\mathbf{q}) = (2\pi)^3 n_e (2\pi m_e k_B T_e)^{-3/2} \exp\left(-\frac{q^2}{2m_e k_B T_e}\right)$, and the non Maxwellian part, $\Delta g(\mathbf{q})$, where $\mathbf{q}$ is a momentum of electron, and n_e , T_e and m_e are electron number density, temperature and mass, respectively. The

heat current density, $\kappa \nabla T_e$, is described by the $\Delta g(\mathbf{q})$ as

$$\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{2m_e} q^2 \frac{\mathbf{q}}{m_e} \Delta g(\mathbf{q}) = -f_\kappa \kappa_{\text{Sp}} \nabla T_e,$$

where f_κ is the heat conductivity in the Spitzer value unit and κ_{Sp} is the Spitzer heat conductivity. The Spitzer heat conductivity is given by $\kappa_{\text{Sp}} = 1.31 \lambda_e n_e k_B \left(\frac{k_B T_e}{m_e} \right)^{1/2}$ where λ_e is the electron Coulomb mean free path (Spitzer 1956; Sarazin 1988). The deviation of the electron momentum distribution from the thermal equilibrium distribution (Okabe & Hattori 2003) in the plasma with a temperature gradient is described by

$$\Delta g(\mathbf{q}) = f_\kappa \kappa_{\text{Sp}} \frac{m}{k_B n_e} \frac{\mathbf{q} \cdot \nabla T_e}{q_{th}^2 T_e} \left(2 - \frac{4q^2}{5q_{th}^2} \right) g_m, \quad (1)$$

where $q_{th} = \sqrt{2m_e k_B T_e}$ is the electron thermal momentum. The plasma instability is derived by this local deviation of the electron momentum distribution from the thermal equilibrium distribution. It has been shown that the instability may provide an origin of the magnetic fields in cluster of galaxies (Okabe & Hattori 2003). Therefore, the measurement of the functional form of the non-equilibrium momentum distribution function is crucial for confirming the above mentioned theoretical models on the suppression of the heat conductivities and generation of the magnetic fields.

In this paper, a spectrum distortion of the cosmic microwave background (hereafter CMB) due to the inverse Compton scattering with electrons which support heat conduction, is examined by solving the Boltzmann equation for Compton scattering to third-order in the electron momentum. The result provides a direct method for measuring the heat conductivities and the functional form of the electron momentum distribution functions in the plasma with a temperature gradient.

2. The Boltzmann equation for third-order Compton scattering

The spectrum distortion of the cosmic microwave background (CMB) due to the Compton scattering from electrons with momentum distribution function of $g(\mathbf{q})$ is given by the Boltzmann equation (Hu et al. 1994) as

$$\begin{aligned} \frac{Df(\mathbf{x}, \mathbf{p})}{cDt} &= \frac{1}{2E(\mathbf{p})} \int D\mathbf{q} D\mathbf{q}' D\mathbf{p}' (2\pi)^4 \delta^{(4)}(\vec{p} + \vec{q} - \vec{p}' - \vec{q}') |M|^2 \\ &\times \{g(\mathbf{x}, \mathbf{q}') f(\mathbf{x}, \mathbf{p}') [1 + f(\mathbf{x}, \mathbf{p})] - g(\mathbf{x}, \mathbf{q}) f(\mathbf{x}, \mathbf{p}) [1 + f(\mathbf{x}, \mathbf{p}')] \}, \end{aligned} \quad (2)$$

where $f(\mathbf{x}, \mathbf{p})$ is the photon occupation number with a three-momentum $\mathbf{p}$, $\mathbf{p}$ and $\mathbf{p}'$ are the three-momentum for photons of four-momentum $\vec{p}$ and $\vec{p}'$, $\mathbf{q}$ and $\mathbf{q}'$ are the three-momentum

for electrons of four-momentum $\vec{q}$ and $\vec{q}'$, $E(\mathbf{p})$ is the energy of the photon with a momentum $\mathbf{p}$, $|M|^2$ is the Lorentz-invariant matrix element for Compton scattering (Hu et al. 1994), and

$$D\mathbf{q} = \frac{d^3\mathbf{q}}{(2\pi)^3 2E(\mathbf{q})}$$

is the Lorentz-invariant phase space volume element of electrons with an energy of $E(\mathbf{q})$ and a momentum $\mathbf{q}$. The left hand side of the equation is the total derivative of the photon distribution function in the phase space written as,

$$\frac{Df}{Dt} = \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x^i} \frac{dx^i}{dt} + \frac{\partial f}{\partial n_i} \frac{dn_i}{dt} + \frac{\partial f}{\partial E(p)} \frac{dE(p)}{dt},$$

where n_i is the i -th component of the direction cosines, $\mathbf{n}$, for a photon of four-momentum $\vec{p}$. The first two terms in the right hand side is a derivative along the photon path. The third term represents an effect of gravitational deflect of the photon path due to the cluster gravitational lensing effect. Since the deflection angle due to the cluster gravitational lensing effect is at most $30''$, the change of the path length due to the lensing effect is at most 0.01pc and negligibly small compared with the cluster size of $\sim 1\text{Mpc}$. The last term represents the cosmological redshift and the energy change of the photons due to the gravitational scattering by the tangential peculiar motion of the cluster (Birkinshaw et al. 1984; Aso et al. 2002). The gravitational scattering causes the CMB temperature fluctuation of $\delta T/T \sim 10^{-7}$ (Aso et al. 2002). The effect is order of magnitude smaller than the Compton scattering effect concerned in this paper as shown in the next section. Therefore, the terms representing the gravitational lensing effect and the gravitational scattering effect are neglected in the course of the following calculations. Further, the cosmological redshift effect is neglected for simplicity since the effect is trivial (Hu et al. 1994).

The right hand side of the Boltzmann equation is the collision term. The first and the second terms are the contributions from scattering into and out of the momentum state $\mathbf{p}$ including the stimulated emission effects, respectively. In the rest frame of the incident electron, the matrix element for Compton scattering summed over polarization is given by,

$$|M|^2 = 2(4\pi)^2 e^2 \left[\frac{\tilde{p}'}{\tilde{p}} + \frac{\tilde{p}}{\tilde{p}'} - 1 + (\tilde{\mathbf{n}} \cdot \tilde{\mathbf{n}}')^2 \right],$$

where e is the absolute value of the charge of the electron. The variables with $\sim$ denotes those in the electron rest frame, otherwise those in the laboratory frame.

As an example, the case when the electron velocity distribution function is the Maxwell-Boltzmann and the thermal energy contained in electron system is much less than the electron rest mass energy, is considered. By expanding the collision term of the Boltzmann equation

for isotropically distributed incident photons up to the second-order in electron momentum, the Kompaneets equation is deduced (Hu et al. 1994; Kompaneets 1957). The spectrum distortion of the CMB due to the Compton scattering from thermal electrons contained in the clusters of galaxies are obtained by solving the Kompaneets equation under the conditions that the probability of the multiple scattering of a photon is negligibly small and the electron temperature is much higher than the temperature of the CMB. This effect is known as the thermal Sunyaev-Zel'dovich effect (hereafter thermal SZE) (Sunyaev & Zel'dovich 1972).

The spectrum distortion due to the plasma with a temperature gradient is deduced by the following way. In the plasma with a temperature gradient, the electron momentum distribution function is described as $g(\mathbf{q}) = g_m(\mathbf{q}) + \Delta g(\mathbf{q})$. Since the multiple scattering of a photon is negligible in the clusters of galaxies, the effects of the spectrum distortion of the CMB due to the Compton scattering from electrons belonging to $g_m(\mathbf{q})$ and electrons belonging to $\Delta g(\mathbf{q})$ are decoupled. Therefore, each effects can be evaluated separately and the total spectrum distortion is given by the superposition of each results. Hereafter, the SZE due to electrons belonging to $\Delta g(\mathbf{q})$ is referred to gradT SZE. Since the first non-zero moment of the $\Delta g(\mathbf{q})$ appears from the third-order moment, to evaluate the effect of the $\Delta g(\mathbf{q})$ on the CMB spectrum distortion the third order expansion in electron momentum of the collision term of the Boltzmann equation is required. The technical details are summarized in Appendix. A final result of third-order expansion is reduced to

$$\begin{aligned} \frac{Df(\mathbf{x}, \mathbf{p})}{cDt} &= 3f_\kappa \kappa_{\text{Sp}} \frac{\sigma_T}{mc^3} \mathbf{n} \cdot \nabla T_e \\ &\times \left[x f'(\mathbf{x}, \mathbf{p}) + \frac{94}{75} x^2 f''(\mathbf{x}, \mathbf{p}) + \frac{14}{75} x^3 f'''(\mathbf{x}, \mathbf{p}) \right], \end{aligned} \quad (3)$$

where $x = \frac{h\nu}{k_B T_\gamma}$, h is the Planck constant, c is a speed of light, ν is the frequency of the CMB photon, T_γ is the temperature of the CMB, σ_T is the Thomson cross section and $\mathbf{n} \equiv \frac{\mathbf{p}}{p}$ is the direction cosine of the observed photon, respectively. Although the resultant photon distribution is anisotropic, the isotropic photon distribution for incident photons is assumed in the course of calculation since the multiple scattering is negligible and the incident photons are CMB photons. The quantities $k_B T_e / mc^2$ and $E(\mathbf{p}) / mc^2$ are treated as the order of $(v/c)^2$ variables.

3. The spectrum of the gradient-T SZE

The spectrum of the gradT SZE is obtained by inserting the Planck function in the right hand side of the above obtained third-order Boltzmann equation and integrating over line of sight. The frequency dependent amplitude of the distortion expressed in temperature is

obtained as

$$\Delta T_{\nu}^{\nabla T} = \frac{12\sigma_T}{mc^3} T_{\gamma} \int d\ell f_{\kappa} \kappa_{\text{Sp}} \mathbf{n} \cdot \nabla T_e \frac{1}{x^2} \Delta^{\nabla T}(x), \quad (4)$$

$$\Delta^{\nabla T}(x) \equiv -\frac{x^4 e^x}{(e^x - 1)^2} \left[1 - \frac{94}{75} x \coth \frac{x}{2} + \frac{14}{75} x^2 \left(\coth^2 \frac{x}{2} + \frac{1}{2 \sinh^2 \frac{x}{2}} \right) \right], \quad (5)$$

where $d\ell$ is the line of sight integral. Figure 1 compares the spectrum shape of the gradT SZE with the thermal and kinetic SZE(Hu et al. 1994; Birkinshaw 1999). Figure 2 shows the frequency dependence of the kernel function $\Delta^{\nabla T}(x)$ to see diagnostics of the spectrum shape of the gradT SZE compared with the thermal and kinetic SZE(Hu et al. 1994; Birkinshaw 1999). The spectrum shows distinct characteristics compared with others. Therefore, the functional form of the non-Maxwellian part can be measured by this spectrum shape of the gradT SZE. It has a finite value at the cross over frequency of the thermal SZE, that is 218GHz. The location of maxima, cross over frequency, and minima are independent of T_e . They are at $x_{\text{max}} = 3.44$ ($\nu = 195\text{GHz}$), $x_{\text{zero}} = 5.75$ ($\nu = 326\text{GHz}$), and $x_{\text{min}} = 8.36$ ($\nu = 437\text{GHz}$), respectively. Since the amplitude of the spectral distortion is proportional to the inner product of the $\mathbf{n}$ with ∇T_e , the sign of the spectrum depends on the relative direction of the line-of-sight to the temperature gradient. When the hotter region locates closer to the observer, the amplitude of the distorted spectrum is larger than the CMB in the Rayleigh-Jeans part. A spherically symmetric system does not show the spectrum distortion due to the gradT SZE since both of increment and decrement with the same amount appear in a single line-of-sight and these are cancelled out. When the line of sight is exactly perpendicular to the temperature gradient, there is no gradT SZE effect.

The amplitude of the spectrum distortion due to the gradT SZE at the peak frequency is evaluated by the following way. The line of sight integral in the right hand side of the equation is reduced to $T_e^{7/2}$. It is independent from the electron density because the heat conductivity does not depend on the electron density explicitly unless it is saturated(Cowie & McKee 1977). As decreasing the electron density the Coulomb mean free path becomes longer. In the region where the electron density is lower than 10^{-5}cm^{-3} with $k_B T_e = 10\text{keV}$, the mean free path is longer than 2Mpc(Sarazin 1988). For the typical rich cluster, the electron density gets lower than 10^{-5}cm^{-3} in the region of 2Mpc apart from the cluster center. Therefore, in such a cluster out skirt the Coulomb mean free path is longer than temperature gradient scales and the heat conduction is saturated. As modeled by the Cowie and Mackee (1977), the saturated heat conductivity could be evaluated by the collisionless diffusion of the hot electrons with the Maxwellian velocity distribution. It is reasonable to assume that the electron velocity distribution is the Maxwellian in the cluster out skirt region where the heat conduction is saturated since the deviation of the velocity distribution function from the thermal equilibrium form is hard to be established without

any collision between particles. Therefore, we assume that $\Delta g = 0$ in the cluster out skirt region where the heat conduction is saturated. For simplicity, we assume that Δg disappears suddenly at the points where the electron Coulomb mean free path becomes as long as the temperature gradient scale. The temperature of these transition points for the closer side and far side to the observer along the line-of-sight are denoted by T_{e2} and T_{e1} , respectively. Then, the amplitude of the gradT SZE is obtained as

$$\Delta T_{\nu}^{\nabla T} = 0.8 \left(\frac{f_{\kappa}}{1.0} \right) \left[\frac{(k_B T_{e2})^{3.5} - (k_B T_{e1})^{3.5}}{(10\text{keV})^{3.5} - (5\text{keV})^{3.5}} \right] \mu\text{K}. \quad (6)$$

The result shows that the amplitude of the gradT SZE is proportional to the heat conductivity. To extract the heat conductivity from the observed amplitude of the gradT SZE, the electron temperature in the cluster out skirt at the both side of the line-of-sight have to be measured. Since it is practically difficult, some assumptions must be made to extract the heat conductivity from the observed temperature map. The expected amplitude is two order of magnitude less than the thermal SZE and an order of magnitude less than the kinetic SZE but order of magnitude larger than the gravitational scattering effect. To distinguish the gradT SZE signal from other SZE signals, measurement of the spectrum by broad band or multi-band observations are required.

4. Applications to clusters of galaxies

The applications of the gradT SZE to three different situations in clusters of galaxies are considered in this section. First case is that the plasma has a smooth distribution of density and temperature. Second and third cases contain one discontinuity, such as a shock and a cold front along the line-of-sight, respectively. An idealized line-of-sight passing through clusters of galaxies is illustrated in Figure 3. The region denoted by D is a discontinuity. The regions denoted by 1 and 2 are smooth regions. The outer edges of these regions are defined by the place where the heat conduction starts to be saturated (Cowie & McKee 1977; Sarazin 1988). For simplicity, the reduction rate of the heat conduction in these smooth regions, f_{κ} , is assumed to be constant through out the regions and suddenly becomes zero at each edges. The electron temperature at the edge of the discontinuity and at the outer edge of each smooth regions are denoted by suffixes D and o , respectively. Since the scale lengths of the discontinuities, L_D , are comparable to or shorter than the Coulomb mean free path, the reduction rate of the heat conduction in the discontinuities, f'_{κ} , may be different from that in the smooth region and could be much less than 1.

The general expression for the amplitude of the gradT SZE for the observer illustrated

in Figure 3 is obtained by performing the line-of-sight integral as

$$\begin{aligned} \Delta T_{\nu}^{\nabla T} \propto & f_{\kappa} [T_e^o(2)^{7/2} - T_e^D(2)^{7/2}] \\ & + f'_{\kappa} [T_e^D(2)^{7/2} - T_e^D(1)^{7/2}] - f_{\kappa} [T_e^o(1)^{7/2} - T_e^D(1)^{7/2}]. \end{aligned}$$

No discontinuity along a line of sight

When there is no discontinuity along a line of sight, the reduction rate of the heat conductivity is constant through out the line of sight. By setting $f'_{\kappa} = f_{\kappa}$, the amplitude of the gradT SZE at $\nu = 195\text{GHz}$ is reduced to

$$\Delta T_{\nu}^{\nabla T} \sim 0.02 \left(\frac{f_{\kappa}}{0.3} \right) \left[\frac{T_e^o(2)^{7/2} - T_e^o(1)^{7/2}}{(5\text{keV})^{7/2} - (3\text{keV})^{7/2}} \right] \mu\text{K}.$$

The amplitude depends on the temperature at both outer edges of the smooth region and the heat conductivity in the smooth region. The amplitude is insensitive to any local variation of the temperature along line-of-sight. When the temperatures at the both outer edges are close to each other, the amplitude of the gradT SZE becomes zero.

Across a shock front

Although the electron distribution function in the cluster shock waves are quit uncertain, it is likely that the shock is collisionless and the thickness of the shock front is much smaller than the electron mean free path. Therefore, it is expected that the heat conductivity in the shock waves is much less than that in the smooth region, e.g. $f'_{\kappa} \ll f_{\kappa}$. The amplitude of the gradT SZE across a shock front at $\nu = 195\text{GHz}$ is reduced to

$$\Delta T_{\nu}^{\nabla T} \sim -12 \left(\frac{f_{\kappa}}{0.3} \right) \left[\frac{T_e^D(2)^{7/2} - T_e^D(1)^{7/2}}{(30\text{keV})^{7/2} - (5\text{keV})^{7/2}} \right] \mu\text{K},$$

where the effect comes from the difference of the electron temperatures at the both sides of the outer edges in the smooth regions is neglected. The adopted temperature of 30keV as the post shock gas temperature is somewhat larger than the average X-ray temperature of the cluster hot gas, that is 2-10keV. However, there are several observational and theoretical evidences which indicate that the temperature of the shock heated gas due to a cluster-cluster major merger could be as hot as 30keV. A cluster 1E0657-56 is an on-going merging cluster in which a contact discontinuity and a shock are identified (Markevitch et al. 2002). The temperature map of this cluster indicates the existence of the high temperature region hotter than 20keV. The combined analysis of the multi wave band SZE mapping observations of the most X-ray luminous cluster RXJ1347-1145 (Kitayama et al. 2004) showed that the inferred temperature of the south-east part of the cluster is well in excess of 20keV which is confirmed by the Chandra X-ray temperature measurement (Allen et al. 2002). A numerical

simulation of the head on collision of rich clusters (Takizawa 1999) showed that the shock heated gas temperature may temporary exceed 20keV. Therefore, in such an ideal case, the amplitude of the gradT SZE could be comparable to that of the kinetic SZE. An important point we have to emphasize is that the amplitude of the gradT SZE across the shock front is proportional to the heat conductivity in the smooth region. Therefore, this observation can be used to measure the heat conductivity in the smooth region.

Across a cold front

The thickness of cold fronts are yet unresolved. The current observed upper limits are comparable to the electron mean free path. Since the cold fronts are found in relatively large fraction of clusters, the cold fronts should not be a short lived transient structure. To maintain the cold fronts, the heat conductivity should be suppressed in many order of magnitude. It is likely that the heat conductivity in cold fronts is much less than unity, $f'_\kappa \ll 1$, and much less than that in the smooth region, $f'_\kappa \ll f_\kappa$. Therefore, the amplitude of the gradT SZE when the line of sight acrosses one cold front, is estimated as

$$\Delta T_\nu^{\nabla T} \sim -0.1 \left(\frac{f_\kappa}{0.3} \right) \left[\frac{T_e^D(2)^{7/2} - T_e^D(1)^{7/2}}{(8\text{keV})^{7/2} - (4\text{keV})^{7/2}} \right] \mu\text{K},$$

where the effect comes from the difference of the electron temperatures at the both sides of the outer edges in the smooth region is neglected. The dependence on the physical variables are the same as the case acrossing a shock. However, the temperature increase acrossing the cold front is not so significant since there is no thermalization of the kinetic energy as in the case of the shock. Therefore, the expected amplitude is much smaller than the case of acrossing the shock.

5. Discussion

We have shown that the inverse Compton scattering of the CMB photon with electrons in the ICM which has a temperature gradient, results in a new type of the spectrum distortion of the CMB by the third-order perturbation theory of the Compton scattering. The spectrum has an universal shape. The cross over frequency appears at 326GHz. The sign of the spectrum depends on the relative direction of the line-of-sight to the direction of the temperature gradient. When the hotter region locates closer to the observer, the intensity becomes brighter than the CMB spectrum in the frequency regions lower than the cross over frequency. Therefore, the effect is distinguishable from other SZE effects by broad band or multi-bands observations of the SZE. The precise measurement of the spectrum shape of the gradT SZE provides an unique opportunity to measure the non-Maxwellian part of the electron velocity distribution function when the ICM has a temperature gradient. The amplitude

of the spectrum distortion is proportional to the heat conductivity. It provides a possibility for a direct measurement of the heat conductivity. The expected amplitude of the gradT SZE signal is more than two order of magnitude smaller than the thermal SZE. The biggest signal for gradT SZE occurs when the line-of-sight across a shock front. The expected signal for this case is as large as $\sim 30\mu\text{K}$ when $f_\kappa = 1$ at maximum. This is $\sim 10\%$ of the thermal SZE. However, there are variety of the effects which distort the thermal SZE spectra, such as the relativistic correction (Birkinshaw 1999), the existence of the non-thermal electrons (Birkinshaw 1999), the hydrodynamical effect of the cluster merger (Koch 2004) and the turbulent motion giving rised by merger shock (Nagai, Kravtsov & Kosowsky 2003). These can become problematic when using multi-frequency separation techniques.

The gradT SZE does not cause any polarization since the quadrapole moment of the CMB intensity distribution averaged over all the electron momentum is zero. The linear polarization of the Compton scattered photons is generated by the systematic transverse motion of the electron system since it leads a finite quadrapole moment of the CMB intensity distribution in the frame moving with the same velocity as the transverse motion of the electron system (Sunyaev & Zel'dovich 1980). As explained in the Section 1 and appendix, the first and second moments of the electron momentum on $\Delta g(\mathbf{q})$ are zero (Okabe & Hattori 2003). Therefore, the $\Delta g(\mathbf{q})$ does not generate a quadrapole moment of the CMB intensity distribution.

The third order Doppler effect could provide the physical explanation of the spectrum distortion in the gradT SZE although it is not satisfactory enough. The scattering can be treated as almost elastic in the electron rest frame, i.e., $\tilde{p}' = \tilde{p}$. The transformation into the laboratory frame induces a Doppler shift and an energy transfer to the photon amount of δp given as

$$\frac{\delta p}{p} = \frac{1 - \vec{\beta} \cdot \mathbf{n}}{1 - \vec{\beta} \cdot \mathbf{n}'} - 1$$

Averaging over the incoming photon direction $\vec{n}$ which is isotropically distributed, we obtain

$$\left\langle \frac{\delta p}{p} \right\rangle \sim \vec{\beta} \cdot \mathbf{n}' + (\vec{\beta} \cdot \mathbf{n}')^2 + (\vec{\beta} \cdot \mathbf{n}')^3.$$

The averaging over electron velocity with velocity distribution function of Δg results in

$$\left\langle \frac{\delta p}{p} \right\rangle \sim -\frac{6}{5n_e} f_\kappa \kappa_{SP} \frac{\mathbf{n} \cdot \vec{\nabla} T_e}{m_e c^3} \sim \pm 0.14 \left(\frac{\lambda_e}{L} \right) \frac{v_{th}}{c} \frac{4k_B T}{m_e c^2},$$

where λ_e is the Coulomb mean free path of electron, L is the temperature gradient scale length and $v_{th} = \sqrt{2k_B T/m_e}$ is the electron thermal velocity. This is an averaged fractional

energy shift of the photon by a single scattering. When the hot region locates foreground of the cold region along the line of sight, the minus sign in the last equality is taken. The scattered photons lose their energy. Therefore, the original Planck spectrum shifts toward low frequency region contrary to the case of the thermal SZE. This results in the increment in the Rayleigh-Jeans part and the decrement in the Wien part contrary to the thermal SZE. When the cold region locates foreground of the hot region, the plus sign in the last equality is taken. The scattered photons gain the energy. In this case, the original Planck spectrum shifts toward high frequency region. The resultant spectrum distortion is the decrement in the Rayleigh-Jeans part and the increment in the Wien part as same as the case of thermal SZE.

Acknowledgments

The authors would like to thank Y.Fujita, T.Kitayama and N.Sugiyama for their valuable comments. MH would like to thank M.Hoshino for motivating him to do this work in the course of the discussion. The authors would also like to thank anonymous referee who provides constructive and objective comments which were helpful for improving the paper. This work is supported by a Grant-in-Aid for the 21st Century COE Program “Exploring New Science by Bridging Particle-Matter Hierarchy” in Tohoku University, funded by the Ministry of Education, Science, Sports and Culture of Japan and a Grant-in-Aid No.16204010 funded by Japan Society for the Promotion of Science.

Appendix

Some technical details for performing the third order perturbation theory for the Compton scattering are summarized. In the following discussion, $p = E(\mathbf{p})$ and $p' = E(\mathbf{p}')$ are used for simplicity.

The third order expansions of each terms contained in the collision integral are obtained as follows. The expansion to third order of the energy conservation can be treated as an expansion in $\Delta p = E(\mathbf{q}) - E(\mathbf{q}')$ as

$$\begin{aligned} \delta(p + q - p' - q') &= \delta(p' - p - \Delta p) \\ &= \delta(p - p') - \Delta p \frac{\partial}{\partial p'} \delta(p' - p) \\ &\quad + \frac{1}{2} \Delta p^2 \frac{\partial^2}{\partial p'^2} \delta(p' - p) - \frac{1}{6} \Delta p^3 \frac{\partial^3}{\partial p'^3} \delta(p' - p). \end{aligned}$$

Since the electron momentum distribution function does not depends on $\mathbf{q}$ in the first term of the collision integral, first performing the integral by $\mathbf{q}$ and $\mathbf{q}$ is replaced with $\mathbf{q}' + \mathbf{p}' - \mathbf{p}$ by using the momentum conservation. Then, Δp is written as

$$\Delta p = \frac{1}{E(\mathbf{q}') + p'} (\mathbf{p}' \cdot (\mathbf{p}' - \mathbf{p}) + \mathbf{q}' \cdot (\mathbf{p}' - \mathbf{p})).$$

For the second term in the collision integral, the integral by $\mathbf{q}'$ is performed first. The energy difference Δp is written as

$$\Delta p = \frac{-1}{E(\mathbf{q}) + p} (\mathbf{p} \cdot (\mathbf{p} - \mathbf{p}') + \mathbf{q} \cdot (\mathbf{p} - \mathbf{p}')).$$

In the pre-scattering electron rest frame, the energy of the scattered photon is described by the energy of the incident electron as

$$\tilde{p}' = \frac{\tilde{p}}{1 + \frac{\tilde{p}}{m_e c^2} (1 - \mathbf{n} \cdot \mathbf{n}')}.$$

The Doppler formula provides a relation between the photon energy in the rest frame and that in the laboratory frame as

$$\tilde{p} = \gamma p (1 - \boldsymbol{\beta} \cdot \mathbf{n}), \quad \tilde{p}' = \gamma p' (1 - \boldsymbol{\beta} \cdot \mathbf{n}'),$$

where γ is the Lorentz factor and $\boldsymbol{\beta} = \mathbf{q}/mc$ is the incident electron velocity in the laboratory frame relative to the speed of light. The Lorentz invariant relation of $\vec{\tilde{p}} \cdot \vec{\tilde{p}'} = \vec{p} \cdot \vec{p}'$ provides a relation of the photon scattering angle between the rest frame and the laboratory frame as

$$\tilde{\mathbf{n}} \cdot \tilde{\mathbf{n}'} = \frac{-\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}') + (\boldsymbol{\beta} \cdot \mathbf{n})(\boldsymbol{\beta} \cdot \mathbf{n}') + \beta^2(1 - \mathbf{n} \cdot \mathbf{n}') + \mathbf{n} \cdot \mathbf{n}'}{1 - \boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}') + (\boldsymbol{\beta} \cdot \mathbf{n})(\boldsymbol{\beta} \cdot \mathbf{n}')}.$$

Since the ratio of the photon energy to the electron rest mass energy is order of β^2 , the third order expansion of $|M|$ in β is expressed as

$$\begin{aligned} |M|^2 &= 2(4\pi)^2 e^2 (1 + \tilde{\mathbf{n}} \cdot \tilde{\mathbf{n}'}) \\ &= 2(4\pi)^2 e^2 [1 + (\mathbf{n} \cdot \mathbf{n}')^2 - 2\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}') \mathbf{n} \cdot \mathbf{n}' (1 - \mathbf{n} \cdot \mathbf{n}') \\ &\quad + (\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}'))^2 (1 - 3\mathbf{n} \cdot \mathbf{n}') (1 - \mathbf{n} \cdot \mathbf{n}') + 2\beta^2 \mathbf{n} \cdot \mathbf{n}' (1 - \mathbf{n} \cdot \mathbf{n}') \\ &\quad + 2(\boldsymbol{\beta} \cdot \mathbf{n})(\boldsymbol{\beta} \cdot \mathbf{n}') \mathbf{n} \cdot \mathbf{n}' (1 - \mathbf{n} \cdot \mathbf{n}') + 2(\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}'))^3 (1 - 2\mathbf{n} \cdot \mathbf{n}') (1 - \mathbf{n} \cdot \mathbf{n}') \\ &\quad - 2(\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}')) \beta^2 (1 - \mathbf{n} \cdot \mathbf{n}') (1 - 2\mathbf{n} \cdot \mathbf{n}') \\ &\quad - 2(\boldsymbol{\beta} \cdot (\mathbf{n} + \mathbf{n}')) (\boldsymbol{\beta} \cdot \mathbf{n})(\boldsymbol{\beta} \cdot \mathbf{n}') (1 - 3\mathbf{n} \cdot \mathbf{n}') (1 - \mathbf{n} \cdot \mathbf{n}')]. \end{aligned}$$

The third order expansion of the electron energy is $E(\mathbf{q}) \sim mc^2(1 + q^2/2m^2c^2)$.

The 1st and 2nd moments of electron momentum for the Δg are zero, and the 3rd moments are obtained as

$$\begin{aligned} \int \frac{d^3\mathbf{q}}{(2\pi)^3} q_i q_j q_k \Delta g &= -\frac{2}{5} m^2 f_\kappa \kappa_{\text{SP}} \partial_x T_e, & \text{for } i = x, j = k = y \text{ or } z, \\ &= -\frac{6}{5} m^2 f_\kappa \kappa_{\text{SP}} \partial_x T_e, & \text{for } i = j = k = x \\ &= 0, & \text{otherwise} \end{aligned}$$

where the x axis is taken to the direction of the temperature gradient. The non-zero terms in the third order expansion of the collision term are only the third order moment of electron momentum and can be evaluated by the above results. The evaluations of the each terms are straight forward.

REFERENCES

- Allen, S.W., Schmidt, R.W., & Fabian, A.C. 2002, MNRAS, **335**, 256
- Aso, O., Hattori, M. & Futamase, T. 2002, ApJ, **576**, L5
- Birkinshaw, M., Gull, S.F. & Hardebeck, H. 1984, Nature, **309**, 34
- Birkinshaw, M. 1999, Phys.Rep., **310**, 97
- Blanton, E.L., Sarazin, C.L., McNamara, B.R., Wise, M.W. 2001, ApJ, **558**, L15
- Cowie, L.L. & McKee, C.F. 1977, ApJ, **211**, 135
- Ettori, S. & Fabian, A.C. 2000, MNRAS, **317**, L57
- Fabian, A.C. *et al.* 2000, MNRAS, **318**, L65
- Fabian, A.C. *et al.* 2001, MNRAS, **321**, L33
- Fujita, Y. *et al.* 2004, ApJ in press
- Hattori, M. & Umetsu, K. 2000, ApJ, **533**, 84
- Hu, W., Scott, D. & Silk, J. 1994, Phys.Rev.D, **49**, 648
- Ikebe, Y. *et al.* 1999, ApJ, **525**, 58
- Kaastra, J.S. *et al.* 2004, AA, **413**, 415

- Kitayama, T. *et al.* 2004, PASJ, **56**, 17
- Koch, P. 2004, JKAS, bf 37, 471
- Kompaneets, A.S. 1957, Sov.Phys.JETP, **4**, 730
- Levinson, A. & Eichler, D. 1992, ApJ, **387**, 212
- Markevitch, M. *et al.* 2000, ApJ, **541**, 542
- Markevitch, M. *et al.* 2002, ApJ, **567**, L27
- Markevitch, M. *et al.* 2003, ApJ, **586**, L19
- McNamara, B.R. *et al.* 2000, ApJ, *534*, L135
- Nagai, D., Kravtsov, A.V., & Kosowsky, A. 2003, ApJ, *587*, 524
- Okabe, N. & Hattori, M. 2003, ApJ, **599**, 964
- Okabe, N. & Hattori, M. 2004, ApJ, submitted
- Ramani, A. & Laval, G. 1978, Phys.Fluids, **21**, 980
- Sarazin, C. L., X-ray emission from Clusters of Galaxies (Cambridge: Cambridge Univ. Press, 1988)
- Shibata, R., *et al.* 2001, ApJ, **549**, 228
- Spitzer, L.Jr. 1956, Physics of Fully Ionized Gases (New York, Interscience)
- Sunyaev, R.A. & Zel’dovich, Ya.B. 1972, Com.Ap.Sp.Phys., **4**, 173
- Sunyaev, R.A. & Zel’dovich, Ya.B. 1980, MNRAS, **190**, 413
- Takizawa, M. 1999, ApJ, **520**, 514
- Vikhlinin, A., Markevitch, M. & Murray, S.S. 2001, ApJ, **549**, L47
- Watanabe, M., *et al.* 1999, ApJ, bf 527, 80
- Zakamska, N. & Narayan, R. 2003, ApJ, **582**, 162




Fig. 1.— Comparison of the CMB spectrum distortion due to the gradT SZE with the thermal SZE and the kinetic SZE. The absolute amplitude of each effects are compared by log-linear plot. A vertical axis is described in K. A horizontal axis is the frequency of the CMB photons normalized by its temperature. The solid line shows the spectrum of the gradT SZE with amplitude shown in Eq.(6). The dashed line shows the spectrum of the thermal SZE assuming an uniform sphere of radius 0.5Mpc with electron density of 10^{-3}cm^{-3} and electron temperature of 10keV. The dotted line shows the spectrum of the kinetic SZE assuming the same uniform sphere moving with a peculiar velocity of 600km/s.




Fig. 2.— The kernel functions of the three SZE spectrums are shown. The solid line shows the kernel function of the gradT SZE. Dashed line and dotted line are the kernel function of the thermal SZE, $\Delta^T(x) = x^4 e^x / (e^x - 1)^2 (x \coth(x/2) - 4)$, and the kinetic SZE, $\Delta^K(x) = x^4 e^x / (e^x - 1)^2$, respectively (Birkinshaw 1999). A vertical axis is described in dimensionless arbitrary unit.




Fig. 3.— An illustration of the line of sight which passes one discontinuity. The line of sight toward an observer is illustrated by an long arrow. The region denoted by D is a discontinuity. The regions denoted by 1 and 2 are the regions where the temperature and density distributions of the hot gas are smooth. The electron temperatures at the edges of the discontinuity is denoted by the super script D and the electron temperature at the outer edges of the cluster is denoted by the super script o , respectively. The fractional heat conductivity in the smooth region is described by f_{κ} and that in the discontinuity is described by f'_{κ} , respectively.